



# Continuity in the Hurst index of the local times of anisotropic Gaussian random fields

Dongsheng Wu<sup>a,\*</sup>, Yimin Xiao<sup>b</sup>

<sup>a</sup> *Department of Mathematical Sciences, University of Alabama in Huntsville, Huntsville, AL 35899, USA*

<sup>b</sup> *Department of Statistics and Probability, Michigan State University, East Lansing, MI 48824, USA*

Received 26 January 2008; received in revised form 19 August 2008; accepted 2 September 2008

---

## Abstract

Let  $\{X^H(t), t \in \mathbb{R}^N\}$ ,  $H \in (0, 1)^N$  be a family of  $(N, d)$ -anisotropic Gaussian random fields with generalized Hurst indices  $H = (H_1, \dots, H_N) \in (0, 1)^N$ . Under certain general conditions, we prove that the local time of  $\{X^{H^0}(t), t \in \mathbb{R}^N\}$  is jointly continuous whenever  $\sum_{\ell=1}^N 1/H_\ell^0 > d$ . Moreover we show that, when  $H$  approaches  $H^0$ , the law of the local times of  $X^H(t)$  converges weakly [in the space of continuous functions] to that of the local time of  $X^{H^0}$ . The latter theorem generalizes the result of [M. Jolis, N. Viles, Continuity in law with respect to the Hurst parameter of the local time of the fractional Brownian motion, *J. Theoret. Probab.* 20 (2007) 133–152] for one-parameter fractional Brownian motions with values in  $\mathbb{R}$  to a wide class of  $(N, d)$ -Gaussian random fields. The main argument of this paper relies on the recently developed sectorial local nondeterminism for anisotropic Gaussian random fields.

© 2008 Elsevier B.V. All rights reserved.

MSC: 60G15; 60G17; 42C40; 28A80

Keywords: Gaussian random fields; Local times; Convergence in law; Sectorial local nondeterminism

---

## 1. Introduction

Gaussian random fields have been extensively studied in probability theory and applied in many scientific areas including physics, engineering, hydrology, biology, economics, just

---

\* Corresponding author. Tel.: +1 256 824 6676; fax: +1 256 824 6173.

E-mail addresses: [dongsheng.wu@uah.edu](mailto:dongsheng.wu@uah.edu) (D. Wu), [xiao@stt.msu.edu](mailto:xiao@stt.msu.edu) (Y. Xiao).

URLs: <http://webpages.uah.edu/~dw0001> (D. Wu), <http://www.stt.msu.edu/~xiaoyimi> (Y. Xiao).

to mention a few. Since many data sets from various areas such as image processing, hydrology, geostatistics and spatial statistics have anisotropic nature in the sense that they have different geometric and probabilistic characteristics along different directions, many authors have proposed applying anisotropic Gaussian random fields as more realistic models. See, for example, [1–4].

Several classes of anisotropic Gaussian random fields have been introduced and studied for theoretical and application purposes. For example, Kamont [5] introduced fractional Brownian sheets and studied some of their regularity properties. Benassi et al. [6] and Bonami and Estrade [3] considered some anisotropic Gaussian random fields with stationary increments. Anisotropic Gaussian random fields also arise naturally in stochastic partial differential equations (see, e.g., [7–10]), in studying the most visited sites of symmetric Markov processes [11], and as spatial or spatiotemporal models in statistics (e.g., [2,12,13]).

Many of these anisotropic Gaussian random fields are governed by their generalized Hurst indices  $H \in (0, 1)^N$  (see Section 2 for the definition of a generalized Hurst index). People often have to use a statistical estimate of the index  $H$  in practice since the exact value of the index is unknown in general. Therefore, a justification of the use of a model is needed in application with an unknown  $H$ . Motivated by this purpose, Jolis and Viles [14] investigated the continuity in law with respect to the Hurst parameter of the local time of real-valued fractional Brownian motions. They proved that the law of the local times of the fractional Brownian motions with Hurst index  $\alpha$  converges weakly to that of the local time of fractional Brownian motion with Hurst index  $\alpha_0$ , when  $\alpha$  tends to  $\alpha_0$ . However, the method they developed there depends heavily on the one-parameter setting and the explicit covariance structure of fractional Brownian motion. It seems hard to apply the method of Jolis and Viles [14] to Gaussian random fields, where “time” parameters are vectors and their covariance structures are more complicated in general.

The main objective of this paper is to provide a general method for studying the continuity of the laws of the local times of Gaussian random fields. More precisely, we prove that, under some mild conditions, the law (in the space of continuous functions) of the local times of  $(N, d)$ -anisotropic Gaussian random fields with generalized Hurst indices  $H$  converges weakly to that of the local time of an  $(N, d)$ -anisotropic Gaussian field with index  $H^0$ , when  $H$  approaches  $H^0$ . Our result generalizes the result of Jolis and Viles [14] for real-valued fractional Brownian motion to a wide class of  $(N, d)$ -anisotropic Gaussian random fields, including fractional Brownian sheets, anisotropic Gaussian fields with stationary increments and the spatiotemporal models in [12,13]. The main ingredient we use in our proof is the recently developed properties of sectorial local nondeterminism for anisotropic Gaussian random fields, see [15–17].

The rest of this paper is organized as follows. Section 2 states the general condition (i.e., Condition A) on Gaussian random fields under investigation. We show that these conditions are satisfied by several classes of Gaussian random fields which are of importance in theory and/or in applications. In Section 3, we recall the definition of local times of vector fields and prove the existence and joint continuity of the local times of Gaussian random fields satisfying Condition A. The key estimate for this paper is stated as Lemma 3.2. In Section 4, we prove the tightness of the laws of local time  $\{L^H\}$  as  $H$  belongs to a neighborhood of a fixed index  $H^0 \in (0, 1)^N$ . In Section 5, we study the convergence in law of local times of the family of Gaussian random fields satisfying Condition A. Finally, we give the proof of our key lemma, Lemma 3.2, in Section 6.

Throughout this paper, we use  $\langle \cdot, \cdot \rangle$  and  $|\cdot|$  to denote the ordinary scalar product and the Euclidean norm in  $\mathbb{R}^m$  respectively, no matter what the value of the integer  $m$  is. In Section  $i$ , unspecified positive and finite constants  $i$  will be numbered as  $c_{i,1}, c_{i,2}, \dots$ .

## 2. General assumptions and examples

For a fixed vector  $H = (H_1, \dots, H_N) \in (0, 1)^N$ , let  $X_0^H = \{X_0^H(t), t \in \mathbb{R}^N\}$  be a real-valued, centered Gaussian random field with  $X_0^H(0) = 0$  a.s. Denote

$$\sigma^2(s, t; H) = \mathbb{E} \left[ X_0^H(s) - X_0^H(t) \right]^2, \quad s, t \in \mathbb{R}^N. \tag{2.1}$$

Let  $I \subseteq \mathbb{R}^N$  be a closed interval in  $\mathbb{R}^N$ . We say a family of Gaussian random fields  $\{X_0^H, H \in (0, 1)^N\}$  satisfies *Condition A* on  $I$  if the following three conditions hold:

**Condition A1.** For all  $s, t \in I$ ,  $\sigma^2(s, t; H)$  is continuous in  $H \in (0, 1)^N$ .

**Condition A2.** There exist positive continuous functions (in  $H$ )  $c_{1,1}(H), \dots, c_{1,4}(H)$  such that for all  $s, t \in I$

$$c_{1,1}(H) \leq \sigma^2(0, t; H) \leq c_{1,2}(H), \tag{2.2}$$

and

$$c_{1,3}(H) \sum_{\ell=1}^N |s_\ell - t_\ell|^{2H_\ell} \leq \sigma^2(s, t; H) \leq c_{1,4}(H) \sum_{\ell=1}^N |s_\ell - t_\ell|^{2H_\ell}. \tag{2.3}$$

**Condition A3.** There exists a positive continuous function (in  $H$ )  $c_{1,5}(H)$  such that for all integers  $n \geq 1$ , all  $u, t^1, \dots, t^n \in I$ ,

$$\text{Var} \left( X_0^H(u) | X_0^H(t^1), \dots, X_0^H(t^n) \right) \geq c_{1,5}(H) \sum_{\ell=1}^N \min_{0 \leq k \leq n} |u_\ell - t_\ell^k|^{2H_\ell}, \tag{2.4}$$

where  $t_\ell^0 = 0$  for all  $\ell = 1, \dots, N$ .

As in [16], an anisotropic Gaussian random field is said to have the property of sectorial local nondeterminism on  $I$  if **Condition A3** is fulfilled.

Throughout this paper, we will call the vector  $H \in (0, 1)^N$  the (generalized) Hurst index of  $X_0^H$ . Without loss of generality, we may assume that

$$0 < H_1 \leq \dots \leq H_N < 1. \tag{2.5}$$

Let  $X^H = \{X^H(t), t \in \mathbb{R}^N\}$  be an  $(N, d)$ -anisotropic Gaussian random field with Hurst index  $H$  defined by

$$X^H(t) = \left( X_1^H(t), \dots, X_d^H(t) \right), \quad t \in \mathbb{R}^N, \tag{2.6}$$

where  $X_1^H, \dots, X_d^H$  are  $d$  independent copies of  $X_0^H$ . We still say a family of Gaussian random fields  $\{X^H, H \in (0, 1)^N\}$  satisfies *Condition A* on  $I$  if the corresponding real-valued family  $\{X_0^H, H \in (0, 1)^N\}$  satisfies *Condition A* on  $I$ .

Under **Condition A2**, the Gaussian random field  $X^H$  has a version whose sample paths are a.s. continuous on  $I$ . Hence, throughout this paper, we will tacitly assume that the sample paths  $X^H(t)$  are a.s. continuous on  $I$ . For simplicity of notation, later on we will further assume that  $I = [\varepsilon, \varepsilon + 1]^N$ , where  $\varepsilon \in (0, 1)$  is a fixed constant.

For a fixed index  $H \in (0, 1)^N$ , Xiao [16] studied sample path properties of an anisotropic Gaussian random field  $X^H$  satisfying **Conditions A2** and **A3'** (see Eq. (2.15) for a definition of **Condition A3'**), where he established results on the modulus of continuity, small ball probabilities, fractal dimensions, hitting probabilities and local times for  $X^H$ . The emphasis of the present paper is different and we focus on continuity of the laws of the functionals of  $X^H$  as  $H \in (0, 1)^N$  varies.

In the following we provide some important examples of families of Gaussian random fields which satisfy **Condition A**. They cover both isotropic and anisotropic Gaussian random fields, as well as the stationary spatial and spatiotemporal Gaussian models constructed in [12,13].

2.1. Fractional Brownian sheets

For a given vector  $H = (H_1, \dots, H_N) \in (0, 1)^N$ , a real-valued fractional Brownian sheet  $B_0^H = \{B_0^H(t), t \in \mathbb{R}_+^N\}$  with index  $H$  is a centered Gaussian random field with covariance function given by

$$\mathbb{E} \left[ B_0^H(s) B_0^H(t) \right] = \prod_{\ell=1}^N \frac{1}{2} \left( s_\ell^{2H_\ell} + t_\ell^{2H_\ell} - |s_\ell - t_\ell|^{2H_\ell} \right), \quad \forall s, t \in \mathbb{R}_+^N. \tag{2.7}$$

An  $(N, d)$ -fractional Brownian sheet  $B^H = \{B^H(t) : t \in \mathbb{R}_+^N\}$  is defined by

$$B^H(t) = \left( B_1^H(t), \dots, B_d^H(t) \right), \quad \forall t \in \mathbb{R}_+^N, \tag{2.8}$$

where  $B_1^H, \dots, B_d^H$  are  $d$  independent copies of  $B_0^H$ . Due to (2.7),  $B^H$  can be seen as generalizations of one-parameter fractional Brownian motion and the Brownian sheet.

Fractional Brownian sheets have become a typical representative of anisotropic Gaussian random fields since they were first introduced by Kamont [5]. In particular, we believe the methods developed for fractional Brownian sheets can be adapted for studying many spatial and spatiotemporal models with separable covariance structures (see, e.g., [2]).

Many authors have studied the probabilistic, statistical and sample path properties of fractional Brownian sheets. Related to the problems considered in this paper, we mention that Xiao and Zhang [18] and Ayache, Wu and Xiao [19] studied the existence and joint continuity of the local times of fractional Brownian sheet  $B^H$ .

**Proposition 2.1.** *The family of  $(N, d)$ -fractional Brownian sheets  $\{B^H, H \in (0, 1)^N\}$  satisfies **Condition A**.*

**Proof.** Eq. (2.7) implies that for all  $s, t \in I$ ,  $\sigma^2(s, t, H)$  is continuous in  $H \in (0, 1)^N$ . Hence **Condition A1** is satisfied. On the other hand, **Conditions A2** and **A3** follows respectively from the proofs of Lemma 8 in [20] and Theorem 1 in [17]. We omit the details.  $\square$

2.2. Gaussian random fields with stationary increments

Let  $\eta = \{\eta(t), t \in \mathbb{R}^N\}$  be a real-valued centered Gaussian random field with  $\eta(0) = 0$ . We assume that  $\eta$  has stationary increments and continuous covariance function  $R(s, t) = \mathbb{E}[\eta(s)\eta(t)]$ . According to [21],  $R(s, t)$  can be represented as

$$R(s, t) = \int_{\mathbb{R}^N} \left( e^{i\langle s, \lambda \rangle} - 1 \right) \left( e^{-i\langle t, \lambda \rangle} - 1 \right) \Delta(d\lambda) + \langle s, \Sigma t \rangle, \tag{2.9}$$

where  $\Sigma$  is an  $N \times N$  nonnegative definite matrix and  $\Delta(d\lambda)$  is a nonnegative symmetric measure on  $\mathbb{R}^N \setminus \{0\}$  satisfying

$$\int_{\mathbb{R}^N} \frac{|\lambda|^2}{1 + |\lambda|^2} \Delta(d\lambda) < \infty. \tag{2.10}$$

The measure  $\Delta$  and its density (if it exists)  $f(\lambda)$  are called the *spectral measure* and *spectral density* of  $\eta$ , respectively.

It follows from (2.9) that  $\eta$  has stochastic integral representation:

$$\left\{ \eta(t), t \in \mathbb{R}^N \right\} \stackrel{d}{=} \left\{ \int_{\mathbb{R}^N} \left( e^{i\langle t, \lambda \rangle} - 1 \right) W(d\xi) + \langle \mathbf{Y}, t \rangle, t \in \mathbb{R}^N \right\}, \tag{2.11}$$

where  $\stackrel{d}{=}$  denotes equality in all finite-dimensional distributions. In the right-hand side of (2.11),  $\mathbf{Y}$  is a Gaussian random variable with mean 0 and  $W(d\lambda)$  is a centered complex valued Gaussian random measure which is independent of  $\mathbf{Y}$  and satisfies

$$\mathbb{E} \left[ W(A) \overline{W(B)} \right] = \Delta(A \cap B) \quad \text{and} \quad W(-A) = \overline{W(A)}$$

for all Borel sets  $A, B \subseteq \mathbb{R}^N$ . From now on, we will assume  $\mathbf{Y} = 0$ , which is equivalent to assuming  $\Sigma = 0$  in (2.9). Therefore, we have

$$\sigma_\eta^2(h) = \mathbb{E} \left[ (\eta(t+h) - \eta(t))^2 \right] = 2 \int_{\mathbb{R}^N} (1 - \cos\langle h, \lambda \rangle) \Delta(d\lambda). \tag{2.12}$$

Eq. (2.11) provides a useful way for constructing Gaussian random fields with stationary increments by choosing the spectral measure  $\Delta$ . In particular, for  $\alpha \in (0, 1)$ , if  $\Delta$  has a density function  $f$  given by

$$f_\alpha(\lambda) = \frac{1}{|\lambda|^{2\alpha+N}}, \quad \forall \lambda \in \mathbb{R}^N \setminus \{0\}, \tag{2.13}$$

then  $\eta = \{\eta(t), t \in \mathbb{R}^N\}$  is a real-valued fractional Brownian motion of index  $\alpha$ , which is an isotropic Gaussian random field and will be denoted by  $\eta^\alpha$ . Another interesting example of isotropic Gaussian random fields is the fractional Riesz–Bessel motion with indices  $\beta$  and  $\gamma$  introduced by Anh et al. [22], whose spectral density is given by

$$f_{\gamma,\beta}(\lambda) = \frac{1}{|\lambda|^{2\gamma} (1 + |\lambda|^2)^\beta}, \quad \forall \lambda \in \mathbb{R}^N \setminus \{0\}, \tag{2.14}$$

where  $\gamma$  and  $\beta$  are constants satisfying  $\beta + \gamma > \frac{N}{2}$  and  $0 < \gamma < 1 + \frac{N}{2}$ . Anh et al. [22] showed that these Gaussian random fields can be used for modeling simultaneously long range dependence and intermittency; and Xiao [15] studied their sample path properties.

The following example covers a wide class of Gaussian random fields that satisfy Condition A. In fact we will prove a stronger result that these Gaussian random fields satisfy Conditions A1 and A2 and the following.

**Condition A3'.** There exists a positive function  $c_{1,6}(H)$  which is continuous in  $H \in (0, 1)^N$  such that for all integers  $n \geq 1$ , all  $u, t^1, \dots, t^n \in I$ ,

$$\text{Var} \left( X_0^H(u) | X_0^H(t^1), \dots, X_0^H(t^n) \right) \geq c_{1,6}(H) \min_{0 \leq k \leq n} \sum_{\ell=1}^N |u_\ell - t_\ell^k|^{2H_\ell}, \tag{2.15}$$

where  $t^0 = 0$ .

Following [16], an anisotropic Gaussian random field satisfying **Condition A3'** is said to have the property of strong local nondeterminism in the metric  $\rho(s, t) = \sum_{j=1}^N |s_j - t_j|^{H_j}$ . Clearly, **Condition A3'** implies **Condition A3**, but the converse does not hold. Consequently, if a family of anisotropic Gaussian random fields satisfies **Conditions A1, A2 and A3'**, then it satisfies **Condition A**.

**Proposition 2.2.** *Let  $\{\eta^H = \{\eta^H(t), t \in \mathbb{R}^N\}, H = (H_1, \dots, H_N) \in (0, 1)^N\}$  be a family of real-valued centered Gaussian random fields with stationary increments and spectral densities  $\{f(\lambda; H), H \in (0, 1)^N\}$ . Suppose  $f(\lambda; H)$  is continuous in  $H \in (0, 1)^N$  and*

$$f(\lambda; H) \asymp \frac{1}{\left( \sum_{j=1}^N |\lambda_j|^{H_j} \right)^{2+Q}}, \quad \forall \lambda \in \mathbb{R}^N \setminus \{0\}, \tag{2.16}$$

where  $Q = \sum_{\ell=1}^N H_\ell^{-1}$ , and where two functions  $q(t; H) \asymp r(t; H)$  for  $t \in T$  means that there are positive continuous functions  $c_{1,7}(H)$  and  $c_{1,8}(H)$  in  $H$  such that  $c_{1,7}(H) \leq q(t; H)/r(t; H) \leq c_{1,8}(H)$  for all  $t \in T$ . Then the family of Gaussian random fields  $\{\eta^H, H \in (0, 1)^N\}$  satisfies **Conditions A1, A2 and A3'**.

**Proof.** By (2.12), we can write

$$\sigma_{\eta^H}^2(s, t; H) = \sigma_{\eta^H}^2(s - t; H) = 2 \int_{\mathbb{R}^N} (1 - \cos\langle s - t, \lambda \rangle) f(\lambda; H) d\lambda. \tag{2.17}$$

By the continuity of  $f(\lambda; H)$  in  $H$ , (2.16) and the Dominated Convergence Theorem, one can verify that **Condition A1** is satisfied.

In order to verify **Conditions A2 and A3'**, we first derive an appropriate upper bound for  $\sigma_{\eta^H}^2(h; H)$  ( $h \in \mathbb{R}^N$ ) which implies the upper bounds in (2.2) and (2.3), and then prove **Condition A3'**, which also provides the desired lower bounds in (2.2) and (2.3).

Due to (2.16) we may, without of loss of generality, assume  $h_\ell \geq 0$  for all  $\ell = 1, \dots, N$ . By (2.16) and (2.17) and a change of variables  $v_\ell = \left( \sum_{j=1}^N h_j^{H_j} \right)^{H_\ell^{-1}} \lambda_\ell$  ( $\ell = 1, \dots, N$ ), we obtain,

$$\begin{aligned} \sigma_{\eta^H}^2(h) &\leq 2c_{1,8}(H) \int_{\mathbb{R}^N} \frac{1 - \cos\langle h, \lambda \rangle}{\left( \sum_{j=1}^N |\lambda_j|^{H_j} \right)^{2+Q}} d\lambda \\ &= 2c_{1,8}(H) \int_{\mathbb{R}^N} \frac{1 - \cos \left( \sum_{\ell=1}^N \left( \sum_{j=1}^N h_j^{H_j} \right)^{-H_\ell^{-1}} h_\ell v_\ell \right)}{\left( \sum_{j=1}^N |v_j|^{H_j} \right)^{2+Q}} dv \left( \sum_{j=1}^N h_j^{H_j} \right)^2. \end{aligned} \tag{2.18}$$

Since  $\frac{h_\ell^{H_\ell}}{\sum_{j=1}^N h_j^{H_j}} \leq 1$  for all  $\ell = 1, \dots, N$  and the function  $x \mapsto \cos x$  is decreasing in  $(0, \frac{\pi}{2})$ , we derive that

$$\begin{aligned} & \int_{\mathbb{R}^N} \frac{1 - \cos \left( \sum_{\ell=1}^N \left( \sum_{j=1}^N h_j^{H_j} \right)^{-H_\ell^{-1}} h_\ell v_\ell \right)}{\left( \sum_{j=1}^N |v_j|^{H_j} \right)^{2+Q}} dv \\ & \leq \int_{|v| < \pi/2} \frac{1 - \cos \left( \sum_{j=1}^N |v_j| \right)}{\left( \sum_{j=1}^N |v_j|^{H_j} \right)^{2+Q}} dv + \int_{|v| \geq \pi/2} \frac{2}{\left( \sum_{j=1}^N |v_j|^{H_j} \right)^{2+Q}} dv. \end{aligned} \tag{2.19}$$

It can be verified that the last two integrals are convergent (see, e.g., Lemmas 6.3 and 6.4 in [16]). Combining (2.18), (2.19) and the elementary inequality  $\left( \sum_{j=1}^N h_j^{H_j} \right)^2 \leq N \sum_{j=1}^N h_j^{2H_j}$ , we obtain

$$\sigma_{\eta^H}^2(h) \leq c_{1,9}(H) \sum_{j=1}^N h_j^{2H_j}, \tag{2.20}$$

where

$$\begin{aligned} & c_{1,9}(H) \\ & = 2N c_{1,8}(H) \left( \int_{|v| < \pi/2} \frac{1 - \cos \left( \sum_{j=1}^N |v_j| \right)}{\left( \sum_{j=1}^N |v_j|^{H_j} \right)^{2+Q}} dv + \int_{|v| \geq \pi/2} \frac{2}{\left( \sum_{j=1}^N |v_j|^{H_j} \right)^{2+Q}} dv \right), \end{aligned}$$

which is a positive continuous function of  $H \in (0, 1)^N$ . Therefore, the upper bounds in Condition A2 follow from (2.20).

Next, we prove that the family  $\{\eta^H, H \in (0, 1)^N\}$  satisfies Condition A3'. The key technique in our derivation is based on the Fourier analytic argument in [23, Chapter 18]; see [15–17] for further information. By following the proof of Theorem 3.2 in [16] line by line, one can verify that for all integers  $n \geq 1$ , all  $u, t^1, \dots, t^n \in I$ ,

$$\text{Var} \left( \eta^H(u) | \eta^H(t^1), \dots, \eta^H(t^n) \right) \geq c_{1,10}(H) \min_{0 \leq k \leq n} \sum_{\ell=1}^N |u_\ell - t_\ell^k|^{2H_\ell}, \tag{2.21}$$

where  $t^0 = 0$  and  $c_{1,10}(H)$  can be chosen as

$$c_{1,10}(H) := \frac{c}{c_{1,7}(H)} \int_{\mathbb{R}^N} \left( \sum_{j=1}^N |\lambda_j|^{H_j} \right)^{2+Q} |\widehat{\delta}(\lambda)|^2 d\lambda.$$

In the above,  $\widehat{\delta}$  is the Fourier transform of a  $C^\infty(\mathbb{R}^N)$  function  $\delta$  such that  $\delta(0) = 1$  and  $\delta(t) \equiv 0$  for all  $t \in \mathbb{R}^N$  with  $\rho(0, t) = \sum_{j=1}^N |t_j|^{H_j} \geq 1$ . Hence  $\widehat{\delta}(\cdot) \in C^\infty(\mathbb{R}^N)$  as well and  $\widehat{\delta}(\lambda)$  decays rapidly as  $|\lambda| \rightarrow \infty$ . This implies that  $c_{1,10}(H)$  is a positive continuous function in  $H \in (0, 1)^N$ . Consequently, we prove **Condition A3'**.

Finally, we can use the lower bound in **Condition A3'** with  $n = 1$  by choosing  $u = t, t^1 = 0$  and  $u = t, t^1 = s$ , respectively, to serve as the lower bounds in **Condition A2**. This finishes the proof of **Proposition 2.2**.  $\square$

**Remark 2.3.** It follows from (2.13) and (2.14) that the spectral density functions of fractional Brownian motion and fractional Riesz–Bessel motion (with  $0 < \gamma + \beta - \frac{N}{2} < 1$ ) satisfy the spectral conditions in **Proposition 2.2**. Therefore, both families of fractional Brownian motions and fractional Riesz–Bessel motions satisfy **Condition A**. When the index  $\alpha \in (0, 1)$  is fixed, Pitt [24] proved that fractional Brownian motion with Hurst index  $\alpha$  is strongly local nondeterministic, i.e., for all integers  $n \geq 1$ , all  $u, t^1, \dots, t^n \in I$ ,

$$\text{Var} \left( \eta^\alpha(u) | \eta^\alpha(t^1), \dots, \eta^\alpha(t^n) \right) \geq c_{1,11}(\alpha) \min_{0 \leq k \leq n} |u - t^k|^{2\alpha}, \tag{2.22}$$

where  $t^0 = 0$  and  $c_{1,11}(\alpha)$  is a positive constant depending on  $\alpha$ . A similar result was proved by Xiao [15] for fractional Riesz–Bessel motion. However their results do not provide any information on whether the constants in the lower bounds are continuous in the indices  $\alpha, \beta$  and  $\gamma$ . In this sense, **Proposition 2.2** strengthens the results of Pitt [24] and Xiao [15].

**Remark 2.4.** Anisotropic Gaussian random fields with the above type of spectral density functions arise naturally in probability theory and its applications. See [25] for their relevance to the solution of the stochastic heat equation; [3,4,26] for their applications in modeling bone and aquifer structures; [12,13] for stationary nonseparable spatial and spatiotemporal Gaussian models.

### 3. Local times and their joint continuity

In this section, we briefly recall some aspects of the theory of local times in general at first and then study the existence and joint continuity of the local times of Gaussian random fields satisfying **Condition A**. For excellent surveys on local times of random and/or deterministic vector fields, we refer to [27,28].

Let  $Y(t)$  be a Borel vector field on  $\mathbb{R}^N$  with values in  $\mathbb{R}^d$ . For any Borel set  $T \subseteq \mathbb{R}^N$ , the occupation measure of  $Y$  on  $T$  is defined as the following measure on  $\mathbb{R}^d$ :

$$\mu_T(\bullet) = \lambda_N \{t \in T : Y(t) \in \bullet\},$$

where  $\lambda_N$  denotes the Lebesgue measure in  $\mathbb{R}^N$ .

If  $\mu_T$  is absolutely continuous with respect to the Lebesgue measure  $\lambda_d$ , we say that  $Y(t)$  has *local times* on  $T$ , and define its local times,  $L(\bullet, T)$ , as the Radon–Nikodým derivative of  $\mu_T$  with respect to  $\lambda_d$ , i.e.,

$$L(x, T) = \frac{d\mu_T}{d\lambda_d}(x), \quad \forall x \in \mathbb{R}^d.$$

In the above,  $x$  is the so-called *space variable*, and  $T$  is the *time variable*. Note that if  $Y$  has local times on  $T$  then for every Borel set  $S \subseteq T$ ,  $L(x, S)$  also exists.



By standard martingale and monotone class arguments, one can deduce that the local times have a measurable modification that satisfies the following *occupation density formula* (see [27, Theorem 6.4]): For every Borel set  $T \subseteq \mathbb{R}^N$ , and for every measurable function  $f : \mathbb{R}^d \rightarrow \mathbb{R}_+$ ,

$$\int_T f(Y(t)) dt = \int_{\mathbb{R}^d} f(x)L(x, T) dx. \tag{3.1}$$

Suppose we fix a rectangle  $T = \prod_{i=1}^N [a_i, a_i + h_i] \subseteq \mathbb{R}^N$ , where  $a \in \mathbb{R}^N$  and  $h \in \mathbb{R}_+^N$ . If we can choose a version of the local time, still denoted by  $L(x, \prod_{i=1}^N [a_i, a_i + t_i])$ , such that it is a continuous function of  $(x, t_1, \dots, t_N) \in \mathbb{R}^d \times \prod_{i=1}^N [0, h_i]$ ,  $Y$  is said to have a *jointly continuous local time* on  $T$ . When a local time is jointly continuous,  $L(x, \cdot)$  can be extended to be a finite Borel measure supported on the level set

$$Y_T^{-1}(x) = \{t \in T : Y(t) = x\}; \tag{3.2}$$

see [29] for details. This makes local times, besides being interesting on their own right, a useful tool for studying fractal properties of  $Y$ .

It follows from (25.5) and (25.7) in [27] that for all  $x, y \in \mathbb{R}^d$ ,  $T \subseteq \mathbb{R}^N$  a closed interval and all integers  $n \geq 1$ ,

$$\begin{aligned} \mathbb{E} [L(x, T)^n] &= (2\pi)^{-nd} \int_{T^n} \int_{\mathbb{R}^{nd}} \exp \left( -i \sum_{j=1}^n \langle u^j, x \rangle \right) \\ &\quad \times \mathbb{E} \exp \left( i \sum_{j=1}^n \langle u^j, Y(t^j) \rangle \right) d\bar{u} d\bar{t} \end{aligned} \tag{3.3}$$

and for all even integers  $n \geq 2$ ,

$$\begin{aligned} \mathbb{E} [(L(x, T) - L(y, T))^n] &= (2\pi)^{-nd} \int_{T^n} \int_{\mathbb{R}^{nd}} \prod_{j=1}^n \left[ e^{-i\langle u^j, x \rangle} - e^{-i\langle u^j, y \rangle} \right] \\ &\quad \times \mathbb{E} \exp \left( i \sum_{j=1}^n \langle u^j, Y(t^j) \rangle \right) d\bar{u} d\bar{t}, \end{aligned} \tag{3.4}$$

where  $\bar{u} = (u^1, \dots, u^n)$ ,  $\bar{t} = (t^1, \dots, t^n)$ , and each  $u^j \in \mathbb{R}^d$ ,  $t^j \in T$ . In the coordinate notation we then write  $u^j = (u_1^j, \dots, u_d^j)$ .

Let  $\{X^H, H \in (0, 1)^N\} = \{X^H(t), t \in \mathbb{R}^N\}$ ,  $H \in (0, 1)^N$  be a family of  $(N, d)$ -Gaussian random fields. For a fixed index  $H^0 \in (0, 1)^N$  such that  $\sum_{\ell=1}^N \frac{1}{H_\ell^0} > d$ , Xiao [16] proved the following results on the existence and joint continuity of the local times of Gaussian random field  $X^{H^0}$ :

- (i) If  $X^{H^0}$  satisfies **Condition A2** (for  $H^0$ ), then  $X^{H^0}$  has a local time  $L^{H^0}(x, T) \in L^2(\mathbb{P} \times \lambda_d)$ .
- (ii) If  $X^{H^0}$  satisfies **Conditions A2** and **A3'** (for  $H^0$ ), then  $X^{H^0}$  has a jointly continuous local time on  $T$ .

The main result of this section is the following **Theorem 3.1**. It shows that, under **Conditions A2** and **A3** (instead of **A3'**), the above conclusions still hold for all  $H \in (0, 1)^N$  which are close to  $H^0$ . Moreover, we can bound the moments of the local times of  $X^H$  in terms

of  $H^0$ . Hence, **Theorem 3.1** strengthens and extends the results of Ayache, Wu and Xiao [19] for fractional Brownian sheets and Theorem 8.2 in [16].

We set up some notation. Let  $H^0 \in (0, 1)^N$  be an index satisfying

$$\sum_{\ell=1}^N \frac{1}{H_\ell^0} > d. \tag{3.5}$$

With the convention  $\sum_{\ell=1}^0 \frac{1}{H_\ell^0} := 0$ , we can see that there exists an integer  $\tau_0 \in \{1, \dots, N\}$  such that

$$\sum_{\ell=1}^{\tau_0-1} \frac{1}{H_\ell^0} \leq d < \sum_{\ell=1}^{\tau_0} \frac{1}{H_\ell^0}. \tag{3.6}$$

Define

$$\beta_{\tau_0} = \sum_{\ell=1}^{\tau_0} \frac{H_{\tau_0}^0}{H_\ell^0} + N - \tau_0 - H_{\tau_0}^0 d. \tag{3.7}$$

Then it can be easily verified that  $\beta_{\tau_0} \in (N - \tau_0, N - \tau_0 + 1]$ , where  $\beta_{\tau_0} = N - \tau_0 + 1$  if and only if  $\sum_{\ell=1}^{\tau_0-1} \frac{1}{H_\ell^0} = d$ ; and, if  $\tau_0 = N$ , then  $\beta_{\tau_0} = H_N^0 (\sum_{\ell=1}^N \frac{1}{H_\ell^0} - d) > 0$ .

Distinguishing two cases  $\sum_{\ell=1}^{\tau_0-1} \frac{1}{H_\ell^0} < d$  and  $\sum_{\ell=1}^{\tau_0-1} \frac{1}{H_\ell^0} = d$ , we see that we can choose a positive number  $\delta_0 < \min\{H_\ell^0, 1 - H_\ell^0 : 1 \leq \ell \leq N\}$ , which depends on  $d$  and  $H^0$  only, with the following property: For all indices  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0] \subseteq (0, 1)^N$ , there is an integer  $\tau \in \{\tau_0 - 1, \tau_0\}$  such that

$$\sum_{\ell=1}^{\tau-1} \frac{1}{H_\ell} \leq d < \sum_{\ell=1}^{\tau} \frac{1}{H_\ell}. \tag{3.8}$$

Moreover, if we denote

$$\beta_\tau = \sum_{\ell=1}^{\tau} \frac{H_\tau}{H_\ell} + N - \tau - H_\tau d, \tag{3.9}$$

then  $\beta_\tau \in (N - \tau_0, N - \tau_0 + 2]$ . It is useful to note that, even though  $\tau$  varies with  $H$ , its value depends only on  $H^0$  and,  $\beta_\tau$  is always bounded from below and above by positive constants depending only on  $H^0$ . In what follows,  $\delta_0$  and  $\tau_0$  will always be the constants defined above.

**Theorem 3.1.** *Let  $\{X^H, H \in (0, 1)^N\} = \{X^H(t), t \in \mathbb{R}^N, H \in (0, 1)^N\}$  be a family of  $(N, d)$ -Gaussian random fields with Hurst indices  $H$  satisfying **Conditions A2** and **A3** on  $I = [\varepsilon, 1 + \varepsilon]^N$ . Let  $H^0 \in (0, 1)^N$  be a Hurst index satisfying (3.5). Then for every  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ ,  $X^H$  has a local time  $L^H(x, I) \in L^2(\mathbb{P} \times \lambda_d)$ , which admits the following  $L^2$ -representation*

$$L^H(x, I) = (2\pi)^{-d} \int_{\mathbb{R}^d} e^{-i\langle y, x \rangle} \int_I e^{i\langle y, X^H(s) \rangle} ds dy, \quad \forall x \in \mathbb{R}^d. \tag{3.10}$$

Furthermore, the Gaussian random field  $X^H = \{X^H(t), t \in \mathbb{R}^N\}$  has almost surely a jointly continuous local time on  $I$ .

For simplicity of notation, we have assumed that  $I = [\varepsilon, 1 + \varepsilon]^N$ . This does not lead to any loss of generality. We will further denote  $L^H(x, t) := L^H(x, [\varepsilon, \varepsilon + t])$ . By Theorem 3.1,  $L^H(x, t)$  is a continuous function on  $\mathbb{R}^d \times [0, 1]^N$ . Hence we will view  $L^H(x, t)$  as an element in  $C(\mathbb{R}^d \times [0, 1]^N, \mathbb{R})$  and  $L^H(x, \cdot)$  as a finite Borel measure. Here and in what follows, for any integers  $p, q$  and Borel set  $S \subseteq \mathbb{R}^p$ ,  $C(S, \mathbb{R}^q)$  denotes the space of continuous functions from  $S$  to  $\mathbb{R}^q$ , endowed with the topology of uniform convergence on compact subsets of  $S$ .

The proof of Theorem 3.1 is based on the following lemma which extends the inequalities in Lemmas 8.4 and 8.8 of [16]. It will also play an essential rôle in Section 4 for proving the tightness of the laws of the local times of  $\{X^H, H \in (0, 1)^N\}$ .

**Lemma 3.2.** Suppose the assumptions of Theorem 3.1 hold. Then, for all  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ , there exist positive and finite constants  $c_{3,1}$  and  $c_{3,2}$  depending on  $N, d, H^0$  and  $I$  only, such that for all hypercubes  $T = [a, a + \langle r \rangle] \subseteq I$  with side length  $r \in (0, 1)$  the following estimates hold:

- (1) For all  $x \in \mathbb{R}^d$  and all integers  $n \geq 1$ ,

$$\mathbb{E} \left[ L^H(x, T)^n \right] \leq c_{3,1}^n (n!)^{N-\beta_\tau} r^{n\beta_\tau}, \tag{3.11}$$

where  $\beta_\tau$  is defined in (3.9).

- (2) For all  $x, y \in \mathbb{R}^d$  with  $|x - y| \leq 1$ , all even integers  $n \geq 1$  and all  $\gamma \in (0, 1)$  small enough,

$$\mathbb{E} \left[ \left( L^H(x, T) - L^H(y, T) \right)^n \right] \leq c_{3,2}^n (n!)^{N-\beta_\tau+(1+H_\tau)\gamma} |x - y|^{n\gamma} r^{n(\beta_\tau-H_\tau\gamma)}. \tag{3.12}$$

The moment estimates (3.11) and (3.12) are a lot more precise than what we actually need in this paper. We expect that they may be useful for some other purposes. For example, one can apply them to show that, for every fixed  $x \in \mathbb{R}^d$ , there is an event of positive probability (which only depends on  $H^0, N, d$  and  $x$ ) such that the Hausdorff dimension of the level set  $\dim_H \left( (X^H)^{-1}(\{x\}) \cap I \right)$  of  $X^H$  tends to  $\dim_H \left( (X^{H^0})^{-1}(\{x\}) \cap I \right)$  as  $H \rightarrow H^0$ . Note that this result cannot be derived directly from the Hausdorff dimension result for the level set of  $X^H$  in [16, Theorem 7.1], where  $H \in (0, 1)^N$  is fixed.

The proof of Lemma 3.2 makes use of Fourier analytic arguments and the property of sectorial local nondeterminism. We will defer the lengthy proof of Lemma 3.2 to Section 6.

**Proof of Theorem 3.1.** Since (3.8) holds for all  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ , the proof of the first part of Theorem 3.1 (i.e., the existence and (3.10)) is the same as that of Theorem 8.1 in [16] and is omitted.

On the other hand, the proof of the joint continuity of the local time of  $X^H$  is similar to that of Theorem 8.2 in [16] (see also the proof of Theorem 3.1 in [19]). Due to its usefulness for proving the tightness in the next section, we include it here. Observe that for all  $x, y \in \mathbb{R}^d, s, t \in [0, 1]^N$  and all even integers  $n \geq 1$ , we have

$$\mathbb{E} \left[ \left( L^H(x, s) - L^H(y, t) \right)^n \right] \leq 2^{n-1} \left\{ \mathbb{E} \left[ \left( L^H(x, s) - L^H(x, t) \right)^n \right] + \mathbb{E} \left[ \left( L^H(x, t) - L^H(y, t) \right)^n \right] \right\}. \tag{3.13}$$

Since  $L^H(x, \cdot)$  is a finite Borel measure, the difference  $L^H(x, s) - L^H(x, t) = L^H(x, [\varepsilon, \varepsilon + s]) - L^H(x, [\varepsilon, \varepsilon + t])$  can be written as a sum of finite number (only depends on  $N$ ) of terms of the form  $L^H(x, T_j)$ , where each  $T_j$  is a closed subinterval of  $I$  with at least one edge length  $\leq |s - t|$ . By further splitting these intervals into cubes of sides  $\leq |s - t|$ , we can use (3.11) to bound the first term in (3.13). On the other hand, the second term in (3.13) can be dealt with using (3.12) as above. Consequently, there exist some constants  $\gamma \in (0, 1)$  and  $n_0$  such that for all  $x, y \in \mathbb{R}^d, s, t \in [0, 1]^N$  and all even integers  $n \geq n_0$ ,

$$\mathbb{E} \left[ \left( L^H(x, s) - L^H(y, t) \right)^n \right] \leq c_{3,3}^n (|x - y| + |s - t|)^{n\gamma}. \tag{3.14}$$

Therefore the joint continuity of the local times  $L^H(x, t)$  follows from the multiparameter version of Kolmogorov’s continuity theorem (cf. [30]). This finishes the proof.  $\square$

**4. Tightness**

In this section, for any index  $H^0 \in (0, 1)^N$  satisfying (3.5), we prove the tightness of the laws of  $\left\{ L^H(x, t), H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0] \right\}$  in  $C([-D, D]^d \times [0, 1]^N, \mathbb{R})$  for all  $D > 0$ .

For this purpose, we will make use of the following tightness criterion which is a consequence of Corollary 16.9 in [31].

**Lemma 4.1.** *Let  $\{Z^{(p)}, p \geq 1\}$  with  $Z^{(p)} = \{Z^{(p)}(t), t \in \mathbb{R}^M\}$  be a sequence of continuous random fields with values in  $\mathbb{R}^q$ . Assume that  $K \subseteq \mathbb{R}^M$  is a compact interval and  $u \in K$  is a fixed point. If there exist some positive constants  $c_{4,1}, b_1, b_2$  and  $b_3$  such that*

$$\mathbb{E} \left[ \left| Z^{(p)}(u) \right|^{b_1} \right] \leq c_{4,1} \quad \forall p \geq 1 \tag{4.1}$$

and

$$\mathbb{E} \left[ \left| Z^{(p)}(s) - Z^{(p)}(t) \right|^{b_2} \right] \leq c_{4,1} |s - t|^{M+b_3} \quad \forall s, t \in K \text{ and } \forall p \geq 1. \tag{4.2}$$

Then  $\{Z^{(p)}, p \geq 1\}$  is tight in  $C(K, \mathbb{R}^q)$ .

**Proposition 4.2.** *Let  $\{X^H, H \in (0, 1)^N\}$  be a family of  $(N, d)$ -Gaussian random fields satisfying Condition A. Let  $H^0 \in (0, 1)^N$  be a Hurst index satisfying (3.5) and  $\delta_0 > 0$  be the corresponding constant defined before. Then the laws of  $\left\{ L^H(x, t), H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0] \right\}$  in  $C([-D, D]^d \times [0, 1]^N, \mathbb{R})$  are tight for all  $D > 0$ .*

**Proof.** Note that, for all  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0], L^H(0, 0) = 0$  almost surely. Hence, by Lemma 4.1, it is sufficient to prove that there exist positive constants  $c_{4,2}$  and  $\gamma \in (0, 1)$  such that for all even integers  $n$  large and all  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ ,

$$\mathbb{E} \left[ \left( L^H(x, s) - L^H(y, t) \right)^n \right] \leq c_{4,2}^n (|x - y| + |s - t|)^{n\gamma}. \tag{4.3}$$

This is similar to (3.14) and the only difference is that the constants  $c_{4,2}$  and  $\gamma$  are independent of  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ . By (3.13) we only need to verify that the upper bounds appearing in the moment estimates (3.11) and (3.12) can be taken to be independent of the index  $H$  provided  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ .

Recall that, by our choice of the constant  $\delta_0$ ,  $\beta_\tau$  is bounded from below and above by positive constants depending only on  $H^0$ . That is, there exist positive constants  $0 < \beta' < \beta''$  such that  $\beta_\tau \in [\beta', \beta'']$  for all  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ . Hence by Lemma 3.2, we can choose  $\gamma \in (0, 1)$  small enough such that

$$\mathbb{E} \left[ L^H(x, T)^n \right] \leq c_{3,1}^n (n!)^{N-\beta'} r^{n\beta'}, \tag{4.4}$$

and

$$\mathbb{E} \left[ \left( L^H(x, T) - L^H(y, T) \right)^n \right] \leq c_{3,2}^n (n!)^{N-\beta'+(1+H_N^0+\delta_0)\gamma} |x - y|^{n\gamma} r^{n(\beta'-(H_N^0+\delta_0)\gamma)} \tag{4.5}$$

for all intervals  $T = [a, a + \langle r \rangle] \subseteq I$  and all  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ . This finishes the proof of Proposition 4.2.  $\square$

### 5. Convergence in law

In this section, we establish the continuity of the laws of the local times of  $X^H$  in the Hurst index  $H \in (0, 1)^N$ . For this purpose, we will make use of the following result, which is an extension of Proposition 4.2 in [14].

**Proposition 5.1.** *Let  $\{Y_n(t), t \in \mathbb{R}^N\}, n \geq 1$  be a family of  $(N, d)$ -random fields satisfying the following conditions:*

- (1)  $\{Y_n\}$  converges in law to  $Y$  in  $C(I, \mathbb{R}^d)$  as  $n \rightarrow \infty$ .
- (2) Both families  $\{Y_n\}$  and  $Y$  have local times  $L_n$  and  $L$ , which are jointly continuous in  $x$  and  $t$ .
- (3) The family of local times  $L_n$  converges in law to a random field  $Z$  in  $C([-D, D]^d \times [0, 1]^N, \mathbb{R})$  as  $n \rightarrow \infty$ .

Then, for all points  $(x^1, t^1), \dots, (x^m, t^m) \in [-D, D]^d \times [0, 1]^N$ , we have

$$\left( Z(x^1, t^1), \dots, Z(x^m, t^m) \right) \stackrel{\mathcal{L}}{=} \left( L(x^1, t^1), \dots, L(x^m, t^m) \right). \tag{5.1}$$

**Proof.** When  $N = 1$  and  $d = 1$ , this is proved by Jolis and Viles [14]. The key ingredient in their proof is the occupation density formula (3.1). Extending their proof to the multiparameter case is straightforward and is omitted.  $\square$

**Theorem 5.2.** *Let  $\{X^H, H \in (0, 1)^N\}$  be a family of  $(N, d)$ -Gaussian random fields satisfying Condition A and let  $H^0 \in (0, 1)^N$  be a Hurst index satisfying (3.5). Then the family of local times  $\{L^H, H \in (0, 1)^N\}$  of  $\{X^H\}$  converges in law to the local time  $L^{H^0}$  of  $X^{H^0}$  in  $C(\mathbb{R}^d \times [0, 1]^N, \mathbb{R})$  as  $H \rightarrow H^0$ .*

**Proof.** It follows from Proposition 16.6 in [31] that it is sufficient to prove that, for all constants  $D > 0$ , the family of local times  $\{L^H, H \in (0, 1)^N\}$  converge in law to the local time  $L^{H^0}$  in  $C([-D, D]^d \times [0, 1]^N, \mathbb{R})$  as  $H \rightarrow H^0$ .

By Proposition 4.2, we see that the laws of  $\left\{ L^H(x, t), H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0] \right\}$  are tight in  $C([-D, D]^d \times [0, 1]^N, \mathbb{R})$  for all  $D > 0$ . Hence it only remains to prove the convergence of finite-dimensional distributions. This can be done by applying Proposition 5.1.

Take an arbitrary sequence  $\{H^n\} \subset \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$  converging to  $H^0$  as  $n \rightarrow \infty$ . First we verify that, as  $n \rightarrow \infty$ , the sequence  $\{X^{H^n}, n \geq 1\}$  of Gaussian random fields converges in law to  $X^{H^0}$  in  $C(I, \mathbb{R}^d)$ . In fact, **Condition A2** implies that for any fixed point  $u \in I$  and all integers  $m \geq 2$ ,

$$\sup_{n \geq 1} \mathbb{E} \left( \left| X^{H^n}(u) \right|^{2m} \right) \leq c_{5,1}^m \tag{5.2}$$

and

$$\mathbb{E} \left( \left| X^{H^n}(u) - X^{H^n}(v) \right|^{2m} \right) \leq c_{5,2}^m \left( \sum_{\ell=1}^N |u_\ell - v_\ell|^{2H_\ell} \right)^m \quad \forall u, v, \in I. \tag{5.3}$$

Hence **Lemma 4.1** implies that the family of laws of  $\{X^H, H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]\}$  is tight in  $C(I, \mathbb{R}^d)$ . On the other hand, **Condition A1** implies that

$$\lim_{H \rightarrow H^0} \mathbb{E} \left[ X_j^H(u) X_k^H(v) \right] = \mathbb{E} \left[ X_j^{H^0}(u) X_k^{H^0}(v) \right], \quad \forall j, k = 1, \dots, d, \forall u, v \in I, \tag{5.4}$$

which implies the convergence of the finite-dimensional distributions of  $\{X^H, H \in (0, 1)^N\}$  as  $H \rightarrow H^0$ . This verifies **Condition (1)** in **Proposition 5.1**. **Condition (2)** in **Proposition 5.1** follows from **Theorem 3.1**.

If for any sequence  $\{H^n\} \subset \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$  converging to  $H^0$  as  $n \rightarrow \infty$  such that

$$L^{H^n}(x, t) \xrightarrow{\mathcal{L}} Z(x, t) \quad \text{in } C([-D, D]^d \times [0, 1]^N, \mathbb{R}) \text{ as } n \rightarrow \infty, \tag{5.5}$$

for some random field  $Z$ . Then, by **Proposition 5.1**, we have that for all fixed points  $(x^1, t^1), \dots, (x^m, t^m)$ ,

$$\left( Z(x^1, t^1), \dots, Z(x^m, t^m) \right) \stackrel{\mathcal{L}}{=} \left( L^{H^0}(x^1, t^1), \dots, L^{H^0}(x^m, t^m) \right), \tag{5.6}$$

which gives us that  $\mathcal{L}(Z) = \mathcal{L}(L^{H^0})$  in  $C([-D, D]^d \times [0, 1]^N, \mathbb{R})$ . This finishes the proof of **Theorem 5.2**.  $\square$

### 6. Proof of Lemma 3.2

The proof of **Lemma 3.2** follows the same spirit of the proofs of **Lemmas 3.7** and **3.10** of [19], where only fractional Brownian sheets were considered. In order to extend their argument to Gaussian random fields satisfying **Condition A** and to prove that the constants  $c_{3,1}$  and  $c_{3,2}$  are independent of  $H$ , we need to make several modifications and rely completely on the sectorial local nondeterminism **A3**.

We will make use of the following lemmas. Among them, **Lemma 6.1** is essentially due to [32] (see also [33]) and **Lemma 6.2** is from [19]. **Lemma 6.3** is a direct consequence of **Condition A3** and tells us that the Gaussian random field  $X_0^H$  has the one-sided strong local nondeterminism along every direction.

**Lemma 6.1.** *Let  $Z_1, \dots, Z_n$  be mean zero Gaussian variables which are linearly independent, then for any nonnegative Borel function  $g : \mathbb{R} \rightarrow \mathbb{R}_+$ ,*

$$\int_{\mathbb{R}^n} g(v_1) \exp \left[ -\frac{1}{2} \text{Var} \left( \sum_{j=1}^n v_j Z_j \right) \right] dv_1 \cdots dv_n$$

$$= \frac{(2\pi)^{(n-1)/2}}{(\det \text{Cov}(Z_1, \dots, Z_n))^{1/2}} \int_{-\infty}^{\infty} g \left( \frac{v}{\sigma_1} \right) e^{-v^2/2} dv,$$

where  $\sigma_1^2 = \text{Var}(Z_1|Z_2, \dots, Z_n)$  is the conditional variance of  $Z_1$  given  $Z_2, \dots, Z_n$ .

**Lemma 6.2.** For any  $q \in [0, \sum_{\ell=1}^N H_\ell^{-1})$ , let  $\tau \in \{1, \dots, N\}$  be the integer such that

$$\sum_{\ell=1}^{\tau-1} \frac{1}{H_\ell} \leq q < \sum_{\ell=1}^{\tau} \frac{1}{H_\ell} \tag{6.1}$$

with the convention that  $\sum_{\ell=1}^0 \frac{1}{H_\ell} := 0$ . Then there exists a positive constant  $\delta_\tau \leq 1$  depending on  $(H_1, \dots, H_N)$  only such that for every  $\delta \in (0, \delta_\tau)$ , we can find  $\tau$  real numbers  $p_\ell \geq 1$  ( $1 \leq \ell \leq \tau$ ) satisfying the following properties:

$$\sum_{\ell=1}^{\tau} \frac{1}{p_\ell} = 1, \quad \frac{H_\ell q}{p_\ell} < 1, \quad \forall \ell = 1, \dots, \tau \tag{6.2}$$

and

$$(1 - \delta) \sum_{\ell=1}^{\tau} \frac{H_\ell q}{p_\ell} \leq H_\tau q + \tau - \sum_{\ell=1}^{\tau} \frac{H_\tau}{H_\ell}. \tag{6.3}$$

Furthermore, if we denote  $\alpha_\tau := \sum_{\ell=1}^{\tau} \frac{1}{H_\ell} - q > 0$ , then for any positive number  $\rho \in (0, \frac{\alpha_\tau}{2\tau})$ , there exists an  $\ell_0 \in \{1, \dots, \tau\}$  such that

$$\frac{H_{\ell_0} q}{p_{\ell_0}} + 2H_{\ell_0} \rho < 1. \tag{6.4}$$

**Lemma 6.3.** Let  $X_0^H$  be an  $(N, 1)$ -Gaussian random field satisfying Condition A3, and let  $\ell \in \{1, \dots, N\}$  be fixed. For any integer  $n \geq 2$ ,  $t^1, \dots, t^n \in I$  such that

$$t_\ell^1 \leq t_\ell^2 \leq \dots \leq t_\ell^n$$

we have

$$\text{Var} \left( X_0^H(t^n) | X_0^H(t^k) : 1 \leq k \leq n-1 \right) \geq c_{1,5}(H) |t_\ell^n - t_\ell^{n-1}|^{2H_\ell}, \tag{6.5}$$

where  $c_{1,5}(H)$  is the positive continuous function defined in Condition A3.

Lemma 6.4 is a refinement of Lemma 3.6 in [19]. It is important to note that the constant  $c_{6,1}$  in (6.6) is independent of  $\{b_j, 1 \leq j \leq n\}$ .

**Lemma 6.4.** Let  $a > 0$  and  $0 < \underline{b} < \bar{b} < 1$  be given constants. There exists a positive constant  $c_{6,1}$  such that for all integers  $n \geq 1$ , real numbers  $0 < r \leq 1$ ,  $b_j \in [\underline{b}, \bar{b}]$  and an arbitrary  $s_0 \in [0, a/2]$ ,

$$\int_{a \leq s_1 \leq \dots \leq s_n \leq a+r} \prod_{j=1}^n (s_j - s_{j-1})^{-b_j} ds_1 \cdots ds_n \leq c_{6,1}^n (n!) \frac{1}{r} \sum_{j=1}^n b_{j-1} r^{n - \sum_{j=2}^n b_j}. \tag{6.6}$$

Please cite this article in press as: D. Wu, Y. Xiao, Continuity in the Hurst index of the local times of anisotropic Gaussian random fields, Stochastic Processes and their Applications (2008), doi:10.1016/j.spa.2008.09.001

In particular, if  $b_j = \alpha \in [\underline{b}, \bar{b}]$  for all  $j = 1, \dots, n$ , then

$$\int_{a \leq s_1 \leq \dots \leq s_n \leq a+r} \prod_{j=1}^n (s_j - s_{j-1})^{-\alpha} ds_1 \cdots ds_n \leq c_{6,1}^n (n!)^{\alpha-1} r^{n(1-(\frac{1}{n})\alpha)}. \tag{6.7}$$

**Proof.** Clearly, we only need to prove (6.6). By integrating the integral in (6.6) in the order of  $ds_n, ds_{n-1}, \dots, ds_1$ , by using a change of variable in each step to construct Beta functions, and by applying the relationship between Beta and Gamma functions, we derive

$$\begin{aligned} & \int_{a \leq s_1 \leq \dots \leq s_n \leq a+r} \prod_{j=1}^n (s_j - s_{j-1})^{-b_j} ds_1 \cdots ds_n \\ &= \frac{1}{1-b_n} \cdot \frac{\Gamma(2-b_n) \prod_{j=2}^{n-1} \Gamma(1-b_j)}{\Gamma\left(n - \sum_{j=2}^n b_j\right)} \int_a^{a+r} (a+r-s_1)^{n-1-\sum_{j=2}^n b_j} (s_1-s_0)^{-b_1} ds_1 \\ &\leq \frac{\prod_{j=1}^n \Gamma(1-b_j)}{\Gamma\left(n - \sum_{j=2}^n b_j\right)} \left(\frac{a}{2}\right)^{-b_1} \int_a^{a+r} (a+r-s_1)^{n-1-\sum_{j=2}^n b_j} ds_1 \\ &= \frac{\prod_{j=1}^n \Gamma(1-b_j)}{\Gamma\left(1+n - \sum_{j=2}^n b_j\right)} \left(\frac{a}{2}\right)^{-b_1} r^{n-\sum_{j=2}^n b_j}. \end{aligned} \tag{6.8}$$

Since the Gamma function  $\Gamma(x)$  is continuous on  $[1-\bar{b}, 1-\underline{b}]$ , there is a finite constant  $c > 0$  such that  $\Gamma(1-b_j) \leq c$  for all  $1 \leq j \leq n$ . Inequality (6.6) follows from (6.8) and Stirling’s formula. □

Now we are ready to prove Lemma 3.2.

**Proof of Lemma 3.2.** For any Gaussian random field  $X_0^H$  satisfying Condition A3, there exists a positive constant

$$c_{6,2} := \min_{H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]} c_{1,5}(H),$$

which depends on  $H^0$  and  $\delta_0$  only, such that for all integers  $n \geq 1$ , all  $u, t^1, \dots, t^n \in I$ ,

$$\text{Var}\left(X_0^H(u) | X_0^H(t^1), \dots, X_0^H(t^n)\right) \geq c_{6,2} \sum_{\ell=1}^N \min_{0 \leq k \leq n} |u_\ell - t_\ell^k|^{2H_\ell}, \tag{6.9}$$

where  $t^0 = 0$ . Meanwhile, by Lemma 6.3, we have that for  $H \in \prod_{\ell=1}^N [H_\ell^0 - \delta_0, H_\ell^0 + \delta_0]$ ,  $\ell \in \{1, \dots, N\}$  fixed, and for any integer  $n \geq 2$ ,  $t^1, \dots, t^n \in I$  such that

$$t_\ell^1 \leq t_\ell^2 \leq \dots \leq t_\ell^n$$



we have

$$\text{Var}\left(X_0^H(t^n) | X_0^H(t^k) : 1 \leq k \leq n-1\right) \geq c_{6,2} |t_\ell^n - t_\ell^{n-1}|^{2H_\ell}. \tag{6.10}$$

Now we proceed to proving (3.11). We will start with an arbitrary closed interval  $T = \prod_{\ell=1}^N [a_\ell, a_\ell + r_\ell] \subseteq I$ . It follows from (3.3) and the fact that  $X_1^H, \dots, X_d^H$  are independent copies of  $X_0^H$  that for all integers  $n \geq 1$ ,

$$\mathbb{E}[L(x, T)^n] \leq (2\pi)^{-nd} \int_{T^n} \prod_{k=1}^d \left\{ \int_{\mathbb{R}^n} \exp\left[-\frac{1}{2} \text{Var}\left(\sum_{j=1}^n u_k^j X_0^H(t^j)\right)\right] dU_k \right\} d\vec{t}, \tag{6.11}$$

where  $U_k = (u_k^1, \dots, u_k^n) \in \mathbb{R}^n$ . Fix  $k = 1, \dots, d$  and denote the inner integral in (6.11) by  $\mathcal{J}_k$ . Since (3.8) holds, we apply Lemma 6.2 with  $\delta = n^{-1}$  and  $q = d$  to obtain  $\tau$  positive numbers  $p_1, \dots, p_\tau \geq 1$  satisfying (6.2) and (6.3). Then for all points  $t^1, \dots, t^n \in T$  such that  $t_\ell^1, \dots, t_\ell^n$  are all distinct for every  $1 \leq \ell \leq N$  [the set of such points has full  $(nN)$ -dimensional Lebesgue measure] we have

$$\begin{aligned} \mathcal{J}_k &= c_{6,3}^n \left[ \det \text{Cov}\left(X_0^H(t^1), \dots, X_0^H(t^n)\right) \right]^{-\frac{1}{2}} \\ &= c_{6,3}^n \prod_{\ell=1}^\tau \left[ \det \text{Cov}\left(X_0^H(t^1), \dots, X_0^H(t^n)\right) \right]^{-\frac{1}{2p_\ell}}, \end{aligned} \tag{6.12}$$

where the first equality follows from the fact that for any positive definite  $q \times q$  matrix  $\Gamma$ ,

$$\int_{\mathbb{R}^q} \frac{[\det(\Gamma)]^{1/2}}{(2\pi)^{u/2}} \exp\left(-\frac{1}{2} x' \Gamma x\right) dx = 1 \tag{6.13}$$

and the second equality follows from (6.2).

Combining (6.11) and (6.12) yields

$$\mathbb{E}[L(x, T)^n] \leq c_{6,4}^n \int_{T^n} \prod_{\ell=1}^\tau \left[ \det \text{Cov}\left(X_0^H(t^1), \dots, X_0^H(t^n)\right) \right]^{-\frac{d}{2p_\ell}} d\vec{t}. \tag{6.14}$$

To evaluate the integral in (6.14), we will first integrate  $[dt_\ell^1 \dots dt_\ell^n]$  for  $\ell = 1, \dots, \tau$ . To this end, we will make use of the following fact about multivariate normal distributions: For any Gaussian random vector  $(Z_1, \dots, Z_n)$ ,

$$\det \text{Cov}(Z_1, \dots, Z_n) = \text{Var}(Z_1) \prod_{j=2}^n \text{Var}(Z_j | Z_1, \dots, Z_{j-1}). \tag{6.15}$$

By the above fact and (6.10), we can derive that for every  $\ell \in \{1, \dots, \tau\}$  and for all  $t^1, \dots, t^n \in T = \prod_{\ell=1}^N [a_\ell, a_\ell + r_\ell]$  satisfying

$$a_\ell \leq t_\ell^{\pi_\ell(1)} \leq t_\ell^{\pi_\ell(2)} \leq \dots \leq t_\ell^{\pi_\ell(n)} \leq a_\ell + r_\ell \tag{6.16}$$

for some permutation  $\pi_\ell$  of  $\{1, \dots, N\}$ , we have

$$\det \text{Cov}\left(X_0^H(t^1), \dots, X_0^H(t^n)\right) \geq c_{6,5}^n \prod_{j=1}^n \left(t_\ell^{\pi_\ell(j)} - t_\ell^{\pi_\ell(j-1)}\right)^{2H_\ell}, \tag{6.17}$$

Please cite this article in press as: D. Wu, Y. Xiao, Continuity in the Hurst index of the local times of anisotropic Gaussian random fields, *Stochastic Processes and their Applications* (2008), doi:10.1016/j.spa.2008.09.001

where  $t_\ell^{\pi_\ell(0)} := \varepsilon$ , and where  $c_{6,5}$  is a constant depending on  $N, H^0, \delta_0$  and  $I$  only. We have chosen  $\varepsilon < \frac{1}{2} \min\{a_\ell, 1 \leq \ell \leq N\}$  so that Lemma 6.4 is applicable.

It follows from (6.16) and (6.17) that

$$\begin{aligned} & \int_{[a_\ell, a_\ell+r_\ell]^n} \left[ \det \text{Cov}(X_0^H(t^1), \dots, X_0^H(t^n)) \right]^{-\frac{d}{2p_\ell}} dt_\ell^1 \cdots dt_\ell^n \\ & \leq \sum_{\pi_\ell} c^n \int_{a_\ell \leq t_\ell^{\pi_\ell(1)} \leq \dots \leq t_\ell^{\pi_\ell(n)} \leq a_\ell+r_\ell} \prod_{j=1}^n \frac{1}{\left(t_\ell^{\pi_\ell(j)} - t_\ell^{\pi_\ell(j-1)}\right)^{H_\ell d/p_\ell}} dt_\ell^1 \cdots dt_\ell^n \\ & \leq c_{6,6}^n (n!)^{H_\ell d/p_\ell} r_\ell^{n\left(1-(1-\frac{1}{n})H_\ell d/p_\ell\right)}. \end{aligned} \tag{6.18}$$

In the above, the last inequality follows from (6.7).

Combining (6.14) and (6.18), and continuing to integrate  $[dt_\ell^1 \dots dt_\ell^n]$  for  $\ell = \tau + 1, \dots, N$ , we obtain

$$\mathbb{E}[L(x, T)^n] \leq c_{6,7}^n (n!)^{\sum_{\ell=1}^{\tau} H_\ell d/p_\ell} \prod_{\ell=1}^{\tau} r_\ell^{n(1-(1-\frac{1}{n})H_\ell d/p_\ell)} \cdot \prod_{\ell=\tau+1}^N r_\ell^n. \tag{6.19}$$

Now we consider the special case when  $T = [a, a + \langle r \rangle]$ , i.e.  $r_1 = \dots = r_N = r$ . Eqs. (6.19) and (6.3) with  $\delta = n^{-1}$  and  $q = d$  together yield

$$\begin{aligned} \mathbb{E}[L(x, T)^n] & \leq c_{6,8}^n (n!)^{\sum_{\ell=1}^{\tau} H_\ell d/p_\ell} r^{n\left(N-(1-n^{-1})\sum_{\ell=1}^{\tau} H_\ell d/p_\ell\right)} \\ & \leq c_{3,1}^n (n!)^{N-\beta_\tau} r^{n\beta_\tau}. \end{aligned} \tag{6.20}$$

This proves (3.11).

We prove estimate (3.12) next. Let  $\gamma \in (0, 1 \wedge \frac{\alpha_\tau}{2\tau})$  be a constant, depending on  $H^0$  only. Note that by the elementary inequalities

$$|e^{iu} - 1| \leq 2^{1-\gamma} |u|^\gamma \quad \text{for all } u \in \mathbb{R} \tag{6.21}$$

and  $|u + v|^\gamma \leq |u|^\gamma + |v|^\gamma$ , we see that for all  $u^1, \dots, u^n, x, y \in \mathbb{R}^d$ ,

$$\prod_{j=1}^n \left| e^{-i\langle u^j, x \rangle} - e^{-i\langle u^j, y \rangle} \right| \leq 2^{(1-\gamma)n} |x - y|^{n\gamma} \sum' \prod_{j=1}^n |u_{k_j}^j|^\gamma, \tag{6.22}$$

where the summation  $\sum'$  is taken over all the sequences  $(k_1, \dots, k_n) \in \{1, \dots, d\}^n$ .

It follows from (3.4) and (6.22) that for every even integer  $n \geq 2$ ,

$$\begin{aligned} & \mathbb{E}[(L(x, T) - L(y, T))^n] \leq (2\pi)^{-nd} 2^{(1-\gamma)n} |x - y|^{n\gamma} \\ & \quad \times \sum' \int_{T^n} \int_{\mathbb{R}^{nd}} \prod_{m=1}^n |u_{k_m}^m|^\gamma \mathbb{E} \exp\left(-i \sum_{j=1}^n \langle u^j, X^H(t^j) \rangle\right) d\bar{u} d\bar{t} \\ & \leq c_{6,9}^n |x - y|^{n\gamma} \sum' \int_{T^n} d\bar{t} \\ & \quad \times \prod_{m=1}^n \left\{ \int_{\mathbb{R}^{nd}} |u_{k_m}^m|^{n\gamma} \exp\left[-\frac{1}{2} \text{Var}\left(\sum_{j=1}^n \langle u^j, X^H(t^j) \rangle\right)\right] d\bar{u} \right\}^{1/n}, \end{aligned} \tag{6.23}$$

where the last inequality follows from the generalized Hölder inequality.

Now we fix a vector  $\bar{k} = (k_1, \dots, k_n) \in \{1, \dots, d\}^n$  and  $n$  points  $t^1, \dots, t^n \in T$  such that  $t_\ell^1, \dots, t_\ell^n$  are all distinct for every  $1 \leq \ell \leq N$ . Let  $\mathcal{M} = \mathcal{M}(\bar{k}, \bar{t}, \gamma)$  be defined by

$$\mathcal{M} = \prod_{m=1}^n \left\{ \int_{\mathbb{R}^{nd}} |u_{k_m}^m|^{n\gamma} \exp \left[ -\frac{1}{2} \text{Var} \left( \sum_{j=1}^n \langle u^j, X^H(t^j) \rangle \right) \right] d\bar{u} \right\}^{1/n}. \tag{6.24}$$

Note that the coordinate fields  $X_\ell^H$  ( $1 \leq \ell \leq N$ ) are independent copies of  $X_0^H$ . By **Condition A3**, the random variables  $X_\ell^H(t^j)$  ( $1 \leq \ell \leq N, 1 \leq j \leq n$ ) are linearly independent. Hence **Lemma 6.1** gives

$$\begin{aligned} & \int_{\mathbb{R}^{nd}} |u_{k_m}^m|^{n\gamma} \exp \left[ -\frac{1}{2} \text{Var} \left( \sum_{j=1}^n \langle u^j, X^H(t^j) \rangle \right) \right] d\bar{u} \\ &= \frac{(2\pi)^{(nd-1)/2}}{[\det \text{Cov}(X_0^H(t^1), \dots, X_0^H(t^n))]^{d/2}} \int_{\mathbb{R}} \left( \frac{v}{\sigma_m} \right)^{n\gamma} e^{-\frac{v^2}{2}} dv \\ &\leq \frac{c_{6,10}^n (n!)^\gamma}{[\det \text{Cov}(X_0^H(t^1), \dots, X_0^H(t^n))]^{d/2}} \frac{1}{\sigma_m^{n\gamma}}, \end{aligned} \tag{6.25}$$

where  $\sigma_m^2$  is the conditional variance of  $X_{k_m}^H(t^m)$  given  $X_i^H(t^j)$  ( $i \neq k_m$  or  $i = k_m$  but  $j \neq m$ ), and the last inequality follows from Stirling’s formula.

Combining (6.24) and (6.25) we obtain

$$\mathcal{M} \leq \frac{c_{6,11}^n (n!)^\gamma}{[\det \text{Cov}(X_0^H(t^1), \dots, X_0^H(t^n))]^{d/2}} \prod_{m=1}^n \frac{1}{\sigma_m^\gamma}. \tag{6.26}$$

For  $\delta = 1/n$  and  $q = d$ , let  $p_\ell$  ( $\ell = 1, \dots, \tau$ ) be the constants as in **Lemma 6.2**. Observe that, since  $\gamma \in (0, \frac{\alpha\tau}{2\tau})$ , there exists an  $\ell_0 \in \{1, \dots, \tau\}$  such that

$$\frac{H_{\ell_0} d}{p_{\ell_0}} + 2H_{\ell_0} \gamma < 1. \tag{6.27}$$

It follows from (6.26) and (6.2) that

$$\mathcal{M} \leq c_{6,12}^n (n!)^\gamma \prod_{\ell=1}^{\tau} \frac{1}{[\det \text{Cov}(X_0^H(t^1), \dots, X_0^H(t^n))]^{d/(2p_\ell)}} \prod_{m=1}^n \frac{1}{\sigma_m^\gamma}. \tag{6.28}$$

The second product in (6.28) will be treated as a ‘‘perturbation’’ factor and will be shown to be small when integrated. For this purpose, we use again the independence of the coordinate processes of  $X^H$  and (6.9) [cf. **Condition A3**] to derive

$$\begin{aligned} \sigma_m^2 &= \text{Var} \left( X_{k_m}^H(t^m) | X_{k_m}^H(t^j), j \neq m \right) \\ &\geq c_{6,13}^2 \sum_{\ell=1}^N \min \left\{ |t_\ell^m - t_\ell^j|^{2H_\ell} : j \neq m \right\}. \end{aligned} \tag{6.29}$$

For any  $n$  points  $t^1, \dots, t^n \in T$ , let  $\pi_1, \dots, \pi_N$  be  $N$  permutations of  $\{1, 2, \dots, n\}$  such that for every  $1 \leq \ell \leq N$ ,

$$t_\ell^{\pi_\ell(1)} \leq t_\ell^{\pi_\ell(2)} \leq \dots \leq t_\ell^{\pi_\ell(n)}. \tag{6.30}$$

Then, by (6.29) and (6.30) we have

$$\begin{aligned} \prod_{m=1}^n \frac{1}{\sigma_m^\gamma} &\leq \prod_{m=1}^n \frac{1}{c_{6,14} \sum_{\ell=1}^N \left[ \left( t_\ell^{\pi_\ell(m)} - t_\ell^{\pi_\ell(m-1)} \right) \wedge \left( t_\ell^{\pi_\ell(m+1)} - t_\ell^{\pi_\ell(m)} \right) \right]^{H_\ell \gamma}} \\ &\leq \prod_{m=1}^n \frac{1}{c_{6,14} \left[ \left( t_{\ell_0}^{\pi_{\ell_0}(m)} - t_{\ell_0}^{\pi_{\ell_0}(m-1)} \right) \wedge \left( t_{\ell_0}^{\pi_{\ell_0}(m+1)} - t_{\ell_0}^{\pi_{\ell_0}(m)} \right) \right]^{H_{\ell_0} \gamma}} \\ &\leq c_{6,14}^{-n} \prod_{m=1}^n \frac{1}{\left( t_{\ell_0}^{\pi_{\ell_0}(m)} - t_{\ell_0}^{\pi_{\ell_0}(m-1)} \right)^{q_{\ell_0}^m H_{\ell_0} \gamma}}, \end{aligned} \tag{6.31}$$

for some  $(q_{\ell_0}^1, \dots, q_{\ell_0}^n) \in \{0, 1, 2\}^n$  satisfying  $\sum_{m=1}^n q_{\ell_0}^m = n$  and  $q_{\ell_0}^1 = 0$ . That is, we will only need to consider the contribution of  $\sigma_m$  in the  $\ell_0$ -th direction.

So far we have obtained all the ingredients for bounding the integral in (6.23) and the rest of the proof is quite similar to the proof of (3.11). It follows from (6.28) and (6.31) that

$$\begin{aligned} \int_{T^n} \mathcal{M}(\bar{k}, \bar{t}, \gamma) d\bar{t} &\leq c_{6,15}^n (n!)^\gamma \int_{T^n} \prod_{\ell=1}^\tau \frac{1}{\left[ \det \text{Cov} \left( X_0^H(t^1), \dots, X_0^H(t^n) \right) \right]^{d/(2p_\ell)}} \\ &\quad \times \prod_{m=1}^n \frac{1}{\left( t_{\ell_0}^{\pi_{\ell_0}(m)} - t_{\ell_0}^{\pi_{\ell_0}(m-1)} \right)^{q_{\ell_0}^m H_{\ell_0} \gamma}} d\bar{t}. \end{aligned} \tag{6.32}$$

To evaluate the above integral, we will first integrate  $[dt_\ell^1 \dots dt_\ell^n]$  for every  $\ell = 1, \dots, \tau$ . Let us first consider  $\ell = \ell_0$ . By using (6.17), (6.15), (6.6) and, thanks to (6.27) and the nature of  $q_{\ell_0}^m$ , we see that

$$\begin{aligned} &\int_{[a_{\ell_0}, a_{\ell_0} + r_{\ell_0}]^n} \frac{1}{\left[ \det \text{Cov} \left( X_0^H(t^1), \dots, X_0^H(t^n) \right) \right]^{d/(2p_{\ell_0})}} \\ &\quad \times \prod_{m=1}^n \frac{1}{\left( t_{\ell_0}^{\pi_{\ell_0}(m)} - t_{\ell_0}^{\pi_{\ell_0}(m-1)} \right)^{q_{\ell_0}^m H_{\ell_0} \gamma}} dt_{\ell_0}^1 \dots dt_{\ell_0}^n \\ &\leq \sum_{\pi_{\ell_0}} c_{6,16}^n \int_{a_{\ell_0} \leq t_{\ell_0}^{\pi_{\ell_0}(1)} \leq \dots \leq t_{\ell_0}^{\pi_{\ell_0}(n)} \leq a_{\ell_0} + r_{\ell_0}} \\ &\quad \times \prod_{m=1}^n \left( t_{\ell_0}^{\pi_{\ell_0}(m)} - t_{\ell_0}^{\pi_{\ell_0}(m-1)} \right)^{-\left( H_{\ell_0} d/p_{\ell_0} + q_{\ell_0}^m H_{\ell_0} \gamma \right)} dt_{\ell_0}^1 \dots dt_{\ell_0}^n \\ &\leq c_{6,17}^n (n!)^{H_{\ell_0} d/p_{\ell_0} + H_{\ell_0} \gamma} r_{\ell_0}^{n \left[ 1 - (1 - \frac{1}{n}) H_{\ell_0} d/p_{\ell_0} - H_{\ell_0} \gamma \right]}. \end{aligned} \tag{6.34}$$

In the above,  $t_{\ell_0}^{\pi_{\ell_0}(0)} = \varepsilon$  as in the proof of (3.11) and the last inequality follows from (6.6).

Meanwhile, recall that, for every  $\ell \neq \ell_0$  ( $\ell \in \{1, \dots, \tau\}$ ), we have shown in (6.18) that

$$\begin{aligned} &\int_{[a_\ell, a_\ell + r_\ell]^n} \left[ \det \text{Cov} \left( X_0^H(t^1), \dots, X_0^H(t^n) \right) \right]^{-\frac{d}{2p_\ell}} dt_\ell^1 \dots dt_\ell^n \\ &\leq c_{6,6}^n (n!)^{H_\ell d/p_\ell} r_\ell^{n \left( 1 - (1 - \frac{1}{n}) H_\ell d/p_\ell \right)}. \end{aligned} \tag{6.35}$$

Finally, we proceed to integrating  $[dt_\ell^1 \dots dt_\ell^n]$  for  $\ell = \tau + 1, \dots, N$ . It follows from the above that

$$\int_{T^n} \mathcal{M}(\bar{k}, \bar{t}, \gamma) d\bar{t} \leq c_{4,22}^n (n!)^{\sum_{\ell=1}^{\tau} H_\ell d/p_\ell + H_{\ell_0}\gamma + \gamma} \times r_{\ell_0}^{n\left[1 - (1 - \frac{1}{n})H_{\ell_0} d/p_{\ell_0} - H_{\ell_0}\gamma\right]} \times \prod_{\ell \neq \ell_0}^{\tau} r_\ell^{n\left[1 - (1 - \frac{1}{n})H_\ell d/p_\ell\right]} \prod_{\ell=\tau+1}^N r_\ell^n. \tag{6.36}$$

In particular, if  $r_1 = \dots = r_N = r \leq 1$ , we combine (6.23) and (6.36) to obtain

$$\begin{aligned} & \mathbb{E} \left[ (L(x, T) - L(y, T))^n \right] \\ & \leq c_{6,18}^n |x - y|^{n\gamma} (n!)^{\sum_{\ell=1}^{\tau} H_\ell d/p_\ell + H_{\ell_0}\gamma + \gamma} \cdot r^{n\left(N - (1 - \frac{1}{n})\sum_{\ell=1}^{\tau} H_\ell d/p_\ell - H_{\ell_0}\gamma\right)} \\ & \leq c_{3,2}^n (n!)^{N - \beta_\tau + (1 + H_\tau)\gamma} |x - y|^{n\gamma} r^{n(\beta_\tau - H_\tau\gamma)}. \end{aligned} \tag{6.37}$$

The last inequality follows from the fact that  $H_{\ell_0} \leq H_\tau$  and Lemma 6.2. This finishes the proof of (3.12). □

**Acknowledgements**

This paper was finished while Y. Xiao was visiting the Statistical & Applied Mathematical Sciences Institute (SAMSI). He thanks the staff of SAMSI for their support and the good working conditions. His research was partially supported by NSF grant DMS-0706728.

The authors thank the referee for his/her helpful suggestions which have led to the improvement of the manuscript.

**References**

- [1] S. Davies, P. Hall, Fractal analysis of surface roughness by using spatial data, *J. Roy. Statist. Soc. Ser. B* 61 (1999) 3–37 (with discussion).
- [2] M.J. Christakos, *Modern Spatiotemporal Geostatistics*, Oxford University Press, 2000.
- [3] A. Bonami, A. Estrade, Anisotropic analysis of some Gaussian models, *J. Fourier Anal. Appl.* 9 (2003) 215–236.
- [4] D.A. Benson, M.M. Meerschaert, B. Baeumer, H.P. Scheffler, Aquifer operator-scaling and the effect on solute mixing and dispersion, *Water Res. Res.* 42 (1) (2006) W01415. 18 pp.
- [5] A. Kamont, On the fractional anisotropic Wiener field, *Probab. Math. Statist.* 16 (1996) 85–98.
- [6] A. Benassi, S. Jaffard, D. Roux, Elliptic Gaussian random processes, *Rev. Mat. Iberoamericana* 13 (1997) 19–90.
- [7] R.C. Dalang, Extending martingale measure stochastic integral with applications to spatially homogeneous s.p.d.e’s, *Electron. J. Probab.* 4 (6) (1999) 1–29.
- [8] B. Øksendal, T. Zhang, Multiparameter fractional Brownian motion and quasi-linear stochastic partial differential equations, *Stochast. Stochast. Rep.* 71 (2000) 141–163.
- [9] C. Mueller, R. Tribe, Hitting probabilities of a random string, *Electron. J. Probab.* 7 (10) (2002) 1–29.
- [10] D. Nualart, Stochastic heat equation driven by fractional noise, 2006, a preprint.
- [11] N. Eisenbaum, D. Khoshnevisan, On the most visited sites of symmetric Markov processes, *Stochastic Process. Appl.* 101 (2002) 241–256.
- [12] T. Gneiting, Nonseparable, stationary covariance functions for space–time data, *J. Amer. Statist. Assoc.* 97 (2002) 590–600.
- [13] M.L. Stein, Space-time covariance functions, *J. Amer. Statist. Assoc.* 100 (2005) 310–321.
- [14] M. Jolis, N. Viles, Continuity in law with respect to the Hurst parameter of the local time of the fractional Brownian motion, *J. Theoret. Probab.* 20 (2007) 133–152.

Please cite this article in press as: D. Wu, Y. Xiao, Continuity in the Hurst index of the local times of anisotropic Gaussian random fields, *Stochastic Processes and their Applications* (2008), doi:10.1016/j.spa.2008.09.001

- [15] Y. Xiao, Strong local nondeterminism and sample path properties of Gaussian random fields, in: Tze Leung Lai, Qiman Shao, Lianfen Qian (Eds.), *Asymptotic Theory in Probability and Statistics with Applications*, Higher Education Press, Beijing, 2007, pp. 136–176.
- [16] Y. Xiao, Sample path properties of anisotropic Gaussian random fields, in: D. Khoshnevisan, F. Rassoul-Agha (Eds.), *A Minicourse on Stochastic Partial Differential Equations*, Springer, New York, 2007 (in press).
- [17] D. Wu, Y. Xiao, Geometric properties of fractional Brownian sheets, *J. Fourier Anal. Appl.* 13 (2007) 1–37.
- [18] Y. Xiao, T. Zhang, Local times of fractional Brownian sheets, *Probab. Theory Related Fields* 124 (2002) 204–226.
- [19] A. Ayache, D. Wu, Y. Xiao, Joint continuity of the local times of fractional Brownian sheets, *Ann. Inst. H. Poincaré Probab. Statist.* 44 (2008) 727–748.
- [20] A. Ayache, Y. Xiao, Asymptotic properties and Hausdorff dimensions of fractional Brownian sheets, *J. Fourier Anal. Appl.* 11 (2005) 407–439.
- [21] A.M. Yaglom, Some classes of random fields in  $n$ -dimensional space, related to stationary random processes, *Theory Probab. Appl.* 2 (1957) 273–320.
- [22] V.V. Anh, J.M. Angulo, M.D. Ruiz-Medina, Possible long-range dependence in fractional random fields, *J. Statist. Plann. Inference* 80 (1999) 95–110.
- [23] J.-P. Kahane, *Some Random Series of Functions*, 2nd ed., Cambridge University Press, Cambridge, 1985.
- [24] L.D. Pitt, Local times for Gaussian vector fields, *Indiana Univ. Math. J.* 27 (1978) 309–330.
- [25] R.S. Robeva, L.D. Pitt, On the equality of sharp and germ  $\sigma$ -fields for Gaussian processes and fields, *Pliska Stud. Math. Bulgar.* 16 (2004) 183–205.
- [26] H. Biermé, M.M. Meerschaert, H.-P. Scheffler, Operator scaling stable random fields, *Stochastic Process. Appl.* 117 (2007) 312–332.
- [27] D. Geman, J. Horowitz, Occupation densities, *Ann. Probab.* 8 (1980) 1–67.
- [28] M. Dozzi, Occupation density and sample path properties of  $N$ -parameter processes, in: *Topics in Spatial Stochastic Processes* (Martina Franca, 2001), in: *Lect. Notes Math.*, vol. 1802, Springer, Berlin, 2002, pp. 127–166.
- [29] R.J. Adler, *The Geometry of Random Fields*, Wiley, New York, 1981.
- [30] D. Khoshnevisan, *Multiparameter Processes: An Introduction to Random Fields*, Springer, New York, 2002.
- [31] O. Kallenberg, *Foundations of Modern Probability*, Springer, New York, 2002.
- [32] J. Cuzick, J. DuPreez, Joint continuity of Gaussian local times, *Ann. Probab.* 10 (1982) 810–817.
- [33] D. Khoshnevisan, Y. Xiao, Images of the Brownian sheet, *Trans. Amer. Math. Soc.* 359 (2007) 3125–3151.